

NCKU-HEP-98-13
hep-ph/9810222

Perturbative QCD analysis of exclusive B meson decays¹

Hsiang-nan Li

Department of Physics, National Cheng-Kung University,
Tainan, Taiwan 701, Republic of China

Abstract

We review the perturbative QCD formalism for exclusive heavy meson decays, concentrating on the three-scale factorization theorem for nonleptonic processes. The formalism is then extended to the radiative decay $B \rightarrow K^*\gamma$, which occurs through penguin diagrams. It is observed that the contributions from the operators other than the penguin one $b \rightarrow s\gamma$ are not negligible. From the best fit to the experimental data of the branching ratio $\mathcal{B}(B \rightarrow K^*\gamma)$, we extract the B meson wave function, which possesses a sharp peak in the region with a small momentum fraction.

¹talk presented at the Fourth Workshop on Particle Physics Phenomenology, Kaohsiung, Taiwan, 1998

1. Introduction

Recently, perturbative QCD (PQCD) has been proposed to be an alternative approach to the study of heavy hadron decays [1, 2, 3], which complements the heavy quark effective theory [4] and the Bauer-Stech-Wirbel (BSW) method [5]. In this talk we review the PQCD analysis of exclusive B meson decays, concentrating on the three-scale factorization theorem for nonleptonic processes [6]. The PQCD formalism is then applied to the radiative decay $B \rightarrow K^* \gamma$, which involves the contributions from penguin diagrams. According to the factorization theorem, a heavy meson decay rate is expressed as the convolution of a hard subamplitude with meson wave functions. The former is calculable in usual perturbation theory, while the latter must be extracted from experimental data or derived by using non-perturbative methods, such as QCD sum rules. Since the K^* meson wave function has been known from the sum rule analyses [7], the $B \rightarrow K^* \gamma$ decay is an ideal process, from which the unknown B meson wave function can be determined. With the B meson wave function obtained here, we are able to make predictions for other B meson decay modes.

The operator for exclusive semileptonic heavy meson decays, such as $B \rightarrow D^{(*)} l \nu$, is

$$\frac{G_F}{\sqrt{2}} V_{cb} \bar{\nu} \gamma_\mu (1 - \gamma_5) l \bar{c} \gamma^\mu b, \quad (1)$$

where G_F is the Fermi coupling constant and V_{cb} the Cabibbo-Kabayashi-Maskawa (CKM) matrix element. The formalism for semileptonic decays is simpler, and part of that for nonleptonic decays. Hence, we shall explain only the factorization of nonleptonic heavy meson decays below, which occur through the Hamiltonian,

$$H = \frac{G_F}{\sqrt{2}} V_{ij} V_{kl}^* (\bar{q}_l q_k) (\bar{q}_j q_i), \quad (2)$$

$(\bar{q}q) = \bar{q} \gamma_\mu (1 - \gamma_5) q$ being the $V - A$ current. Hard gluon corrections cause an operator mixing, and their renormalization-group summation leads to the effective Hamiltonian,

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{ij} V_{kl}^* [c_1(\mu) O_1(\mu) + c_2(\mu) O_2(\mu)], \quad (3)$$

with the four-fermion operators $O_1 = (\bar{q}_l q_k)(\bar{q}_j q_i)$ and $O_2 = (\bar{q}_j q_k)(\bar{q}_l q_i)$. The Wilson coefficients $c_{1,2}$, organizing the large logarithms from the hard gluon corrections to all orders, describe the evolution from the W boson mass M_W to a lower scale μ with the matching conditions $c_1(M_W) = 1$ and $c_2(M_W) = 0$.

The most widely adopted approach to exclusive nonleptonic heavy meson decays is the BSW model [5], in which the factorization hypothesis on the matrix elements of the operators $O_{1,2}$ is assumed. In this model decay rates are expressed in terms of various hadronic transition form factors. Employing the Fierz transformation, the coefficient of the form factors corresponding to external W boson emissions is $a_1 = c_1 + c_2/N$, and that corresponding to internal W boson emissions is $a_2 = c_2 + c_1/N$, N being the number of colors. The form factors may be related to each other by heavy quark symmetry, and parametrized by different ansatz. Nonfactorizable contributions, which can not be expressed in terms of hadronic transition form factors, and nonspectator contributions from W boson exchanges are neglected.

Though the BSW model is simple and gives predictions in fair agreement with experimental data, it encounters several difficulties. It has been known that the large N limit of $a_{1,2}$, *i.e.*, the choice $a_1 = c_1(M_c) \approx 1.26$ and $a_2 = c_2(M_c) \approx -0.52$, with M_c the c quark mass, explains the data of charm decays [5]. However, the same large N limit of $a_1 = c_1(M_b) \approx 1.12$ and $a_2 = c_2(M_b) \approx -0.26$, M_b being the b quark mass, does not apply to the bottom case. That is, the different mechanism between charm and bottom decays can not be understood in the BSW approach. Moreover, it has been difficult to explain the two ratios associated with the $B \rightarrow J/\psi K^{(*)}$ decays [8],

$$R = \frac{\mathcal{B}(B \rightarrow J/\psi K^*)}{\mathcal{B}(B \rightarrow J/\psi K)} , \quad R_L = \frac{\mathcal{B}(B \rightarrow J/\psi K_L^*)}{\mathcal{B}(B \rightarrow J/\psi K^*)} , \quad (4)$$

simultaneously using the BSW model, where $\mathcal{B}(B \rightarrow J/\psi K)$ is the branching ratio of the decay $B \rightarrow J/\psi K$. This controversy was found to be attributed to the neglect of the nonfactorizable internal W -emission amplitudes, which are in fact of the same order as the factorizable ones [9, 10].

The above difficulties have been overcome by a modified PQCD formalism, the three-scale factorization theorem [6]. In the next section we shall summarize the basic ideas of this theorem.

2. Three-scale factorization theorem

Nonleptonic heavy meson decays involve three scales: the W boson mass M_W , at which the matching conditions of the effective Hamiltonian to the original Hamiltonian are defined, the typical scale t of a hard subamplitude, which reflects the dynamics of heavy meson decays, and the factorization scale $1/b$, with b the conjugate variable of parton transverse momenta. The dynamics below $1/b$ is regarded as being completely nonperturbative, and parametrized into a meson wave function $\phi(x)$, x being the momentum fraction. PQCD for the scale above $1/b$ is reliable, and radiative corrections produce two types of large logarithms $\ln(M_W/t)$ and $\ln(tb)$. The former are summed to give the evolution from M_W down to t described by the Wilson coefficients $c(t)$. While the latter are summed to give the evolution from t to $1/b$.

There exist also double logarithms $\ln^2(Pb)$ from the overlap of collinear and soft divergences, P being the dominant light-cone component of a meson momentum. The resummation of these double logarithms leads to a Sudakov form factor $\exp[-s(P, b)]$, which suppresses the long-distance contributions in the large b region, and improves the applicability of PQCD around the energy scale of few GeV. The b quark mass scale is located in the range of applicability. This is the motivation we develop the PQCD formalism for heavy hadron decays. For the detailed derivation of the Sudakov factor for heavy meson decays, refer to [1, 2]. With all the large logarithms organized, the remaining finite contributions are absorbed into a hard b quark decay subamplitude $H(t)$. Because of Sudakov suppression, the perturbative expansion of H in the coupling constant α_s makes sense.

Therefore, a three-scale factorization formula possesses the typical expression,

$$c(t) \otimes H(t) \otimes \phi(x) \otimes \exp \left[-s(P, b) - 2 \int_{1/b}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})) \right], \quad (5)$$

where the exponential containing the quark anomalous dimension γ_q describes the evolution from t to $1/b$ mentioned above. As the hard scale t runs to below M_b and M_c , the constructive and destructive interferences involved in bottom and charm decays, respectively, appear naturally. Furthermore, not only factorizable, but nonfactorizable and nonspectator contributions can be evaluated in a systematic way. With the inclusion of nonfactorizable

contributions, we find that $a_{1,2}$ restore their original role of the Wilson coefficients, instead of being treated as the BSW free parameters. The branching ratios of the various decay modes $B \rightarrow D^{(*)}\pi(\rho)$ and the ratios R and R_L associated with the $B \rightarrow J/\psi K^{(*)}$ decays can all be well explained by the above formalism [11].

We review the evaluation of the factorizable, nonfactorizable and non-spectator contributions to exclusive nonleptonic decays $B \rightarrow D^{(*)}\pi(\rho)$ in the three-scale PQCD factorization theorem. The factorizable external W -emission amplitudes define the $B \rightarrow D^{(*)}$ transition form factors ξ_i , $i = +, -, V, A_1, A_2$, and A_3 , through the hadronic matrix elements,

$$\begin{aligned}\langle D(P_2)|V^\mu|B(P_1)\rangle &= \sqrt{M_B M_D}[\xi_+(\eta)(v_1 + v_2)^\mu + \xi_-(\eta)(v_1 - v_2)^\mu], \\ \langle D^*(P_2)|V^\mu|B(P_1)\rangle &= i\sqrt{M_B M_{D^*}}\xi_V(\eta)\epsilon^{\mu\nu\alpha\beta}\epsilon_\nu^*v_{2\alpha}v_{1\beta}, \\ \langle D^*(P_2)|A^\mu|B(P_1)\rangle &= \sqrt{M_B M_{D^*}}[\xi_{A_1}(\eta)(\eta + 1)\epsilon^{*\mu} - \xi_{A_2}(\eta)\epsilon^* \cdot v_1 v_1^\mu \\ &\quad - \xi_{A_3}(\eta)\epsilon^* \cdot v_1 v_2^\mu].\end{aligned}\tag{6}$$

P_1 (P_2), M_B ($M_{D^{(*)}}$) and v_1 (v_2) are the momentum, the mass, and the velocity of the B ($D^{(*)}$) meson, satisfying the relation $P_1 = M_B v_1$ ($P_2 = M_{D^{(*)}} v_2$). ϵ^* is the polarization vector of the D^* meson. The velocity transfer $v_1 \cdot v_2$ in two-body nonleptonic decays takes the maximal value $\eta = (1 + r^2)/(2r)$ with $r = M_{D^{(*)}}/M_B$. In the infinite mass limit of M_B and $M_{D^{(*)}}$ the form factors ξ_i obey the relations

$$\xi_+ = \xi_V = \xi_{A_1} = \xi_{A_3} = \xi, \quad \xi_- = \xi_{A_2} = 0,\tag{7}$$

where ξ is the so-called Isgur-Wise (IW) function [12].

Equation (6) is the standard definition of the form factors ξ_i . In our approach, however, the factorization formulas for ξ_i will contain the Wilson coefficients as the extra convolution factors. ξ_i include the contributions from the hadronic matrix element of O_1 and from the color-suppressed matrix element of O_2 . Therefore, their factorization formulas involve the Wilson coefficient $a_1 = c_1 + c_2/N$. The form factors for the internal W -emission diagrams include the factorizable contributions from the matrix elements of O_2 and from the color-suppressed matrix element of O_1 , which then contain the Wilson coefficient $a_2 = c_2 + c_1/N$. The form factors for the W -exchange diagrams include the factorizable contributions from the matrix elements of

O_2 and from the color-suppressed matrix element of O_1 . Hence, they also contain the Wilson coefficient a_2 .

The nonfactorizable contributions to the $B \rightarrow D^{(*)}\pi$ decays are color-suppressed, which can not be expressed in terms of hadronic form factors. A nonfactorizable diagram associated with external W emissions involves a hard exchanged gluon, which, for example, attaches the spectator quark of the B meson and a valence quark of the pion. The resultant amplitude depends on the Wilson coefficient c_2/N . A nonfactorizable diagram associated with internal W emissions contains a hard gluon, which attaches the spectator quark of the B meson and a valence quark of the D meson. The resultant amplitude depends on the Wilson coefficient c_1/N . Similarly, a nonfactorizable amplitude associated with W exchanges contains the Wilson coefficient c_1/N . For the explicit expressions of the above form factors and amplitudes, refer to [11].

3. The $B \rightarrow K^*\gamma$ decay

We now apply the three-scale factorization theorem to the radiative decay $B \rightarrow K^*\gamma$ [13], whose effective Hamiltonian is

$$H_{\text{eff}} = -\frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \sum_{i=1}^{12} c_i(\mu) O_i(\mu) . \quad (8)$$

It will be found that only the operators

$$O_2 = (\bar{s}c)(\bar{c}b) , \quad (9)$$

$$O_7 = \frac{e}{8\pi^2} M_b \bar{s} \sigma^{\mu\nu} (1 + \gamma_5) b F_{\mu\nu} , \quad (10)$$

$$O_8 = -\frac{g}{8\pi^2} M_b \bar{s} \sigma^{\mu\nu} (1 + \gamma_5) T^a b G_{\mu\nu}^a , \quad (11)$$

with $\sigma^{\mu\nu} = (i/2)[\gamma^\mu, \gamma^\nu]$, are important. The μ dependence of the corresponding Wilson coefficients $c_2(\mu)$, $c_7(\mu)$ and $c_8(\mu)$ can be quoted from [14].

We assign $P_1 = (M_B/\sqrt{2})(1, 1, \mathbf{0}_T)$ and $P_2 = (M_B/\sqrt{2})(1, r^2, \mathbf{0}_T)$ with $r = M_{K^*}/M_B$, M_{K^*} being the K^* meson mass, as the B and K^* meson momenta, respectively, in the rest frame of the B meson. Assume that the light valence quark in the B (K^*) meson carries the momentum k_1 (k_2). k_1

has a minus component k_1^- , giving the momentum fraction $x_B = k_1^-/P_1^-$, and small transverse components $\mathbf{k}_{BT}$. k_2 has a large plus component k_2^+ , giving $x_K = k_2^+/P_2^+$, and small $\mathbf{k}_{KT}$. The photon momentum is then $P_3 = P_1 - P_2$, whose nonvanishing component is only P_3^- .

The decay amplitude is expressed as

$$M = \epsilon_\gamma \cdot \epsilon_{K^*}^* M_1 + i\epsilon_{\mu\rho+-} \epsilon_\gamma^\mu \epsilon_{K^*}^{*\rho} M_2 , \quad (12)$$

with ϵ_γ and ϵ_{K^*} the polarization vectors of the photon and of the K^* meson, respectively, and

$$M_{1(2)} = M_{1(2)2} + M_{1(2)7} + M_{1(2)8} , \quad (13)$$

where the terms on the right-hand side come from the operators O_2 , O_7 , and O_8 , respectively. From Eq. (12), it is obvious that only the K^* mesons with transverse polarizations are produced in the decay. The factorization formulas for the amplitudes M_{ij} are, in terms of the overall factor,

$$\Gamma^{(0)} = \frac{G_F e}{\sqrt{2} \pi} V_{ts}^* V_{tb} C_F M_B^5 , \quad (14)$$

listed below. The $B \rightarrow K^* \gamma$ decay rate is then given by

$$\Gamma = \frac{1 - r^2}{32\pi M_B} (|M_1|^2 + |M_2|^2) . \quad (15)$$

The amplitude from O_2 is written as

$$\begin{aligned} M_{12} = & \Gamma^{(0)} \frac{4}{3} \int_0^1 dx \int_0^{1-x} dy \int_0^1 dx_B dx_K \int_0^{1/\Lambda} b db \phi_B(x_B) \phi_{K^*}(x_K) \\ & \times \alpha_s(t_2) c_2(t_2) \exp[-S(x_B, x_K, t_2, b, b)] \\ & \times [(1 - r^2 + 2rx_K + 2x_B)y - (rx_K + 3x_B)(1 - x)] \\ & \times \frac{(1 - r)(1 - r^2)x_K x}{xy(1 - r^2)x_K M_B^2 - M_c^2} H_2(Ab, \sqrt{|B_2^2|}b) , \end{aligned} \quad (16)$$

with

$$\begin{aligned} A^2 &= x_K x_B M_B^2 , \\ B_2^2 &= x_K x_B M_B^2 - \frac{y}{1 - x} (1 - r^2) x_K M_B^2 + \frac{M_c^2}{x(1 - x)} , \\ t_2 &= \max(A, \sqrt{|B_2^2|}, 1/b) , \end{aligned} \quad (17)$$

where the integrations over x and y are associated with the charm loop, from which the photon and the hard exchanged gluon are radiated. The variable b_B (b_K), conjugate to the parton transverse momentum k_{BT} (k_{KT}), represents the transverse extent of the B (K^*) meson. t_2 is the characteristic scale of the hard subamplitude,

$$\begin{aligned} H_2(Ab, \sqrt{|B_2^2|}b) &= K_0(Ab) - K_0(\sqrt{|B_2^2|}b), \quad B_2^2 > 0, \\ &= K_0(Ab) - i\frac{\pi}{2}H_0^{(1)}(\sqrt{|B_2^2|}b), \quad B_2^2 < 0, \end{aligned} \quad (18)$$

as mentioned before.

There are two diagrams associated with the operator O_7 , where the hard gluon attaches the two quarks in the B meson and in the K^* meson separately. The corresponding amplitude is

$$\begin{aligned} M_{17} &= \Gamma^{(0)} 2 \int_0^1 dx_B dx_K \int_0^{1/\Lambda} b_B db_B b_K db_K \phi_B(x_B) \phi_{K^*}(x_K) (1 - r^2) \\ &\times \left[r H_7^{(a)}(Ab_K, \sqrt{|B_7^2|}b_B, \sqrt{|B_7^2|}b_K) F_7(t_{7a}) \right. \\ &\left. + [1 + r + (1 - 2r)x_K] H_7^{(b)}(Ab_B, C_7 b_B, C_7 b_K) F_7(t_{7b}) \right], \end{aligned} \quad (19)$$

with

$$\begin{aligned} B_7^2 &= (x_B - r^2) M_B^2, \quad C_7^2 = x_K M_B^2 \\ t_{7a} &= \max(A, \sqrt{|B_7^2|}, 1/b_B, 1/b_K), \\ t_{7b} &= \max(A, C_7, 1/b_B, 1/b_K). \end{aligned} \quad (20)$$

The function F_7 denotes the product

$$F_7(t) = \alpha_s(t) c_7(t) \exp[-S(x_B, x_K, t, b_B, b_K)]. \quad (21)$$

The hard functions

$$\begin{aligned} H_7^{(a)}(Ab_K, \sqrt{|B_7^2|}b_B, \sqrt{|B_7^2|}b_K) &= \\ K_0(Ab_K) h(\sqrt{|B_7^2|}b_B, \sqrt{|B_7^2|}b_K), \quad B_7^2 > 0, \\ K_0(Ab_K) h'(\sqrt{|B_7^2|}b_B, \sqrt{|B_7^2|}b_K), \quad B_7^2 < 0, \end{aligned} \quad (22)$$

with

$$\begin{aligned} h &= \theta(b_B - b_K) K_0(\sqrt{|B_7^2|} b_B) I_0(\sqrt{|B_7^2|} b_K) + (b_B \leftrightarrow b_K) , \\ h' &= i \frac{\pi}{2} \left[\theta(b_B - b_K) H_0^{(1)}(\sqrt{|B_7^2|} b_B) J_0(\sqrt{|B_7^2|} b_K) + (b_B \leftrightarrow b_K) \right] , \end{aligned} \quad (23)$$

and

$$H_7^{(b)}(Ab_B, C_7 b_B, C_7 b_K) = K_0(Ab_B) h(C_7 b_B, C_7 b_K) , \quad (24)$$

come from the two diagrams.

Four diagrams are associated with the operator O_8 , where the photon is radiated by each quark in the B and K^* mesons. The corresponding amplitude is

$$\begin{aligned} M_{18} &= -\Gamma^{(0)} \frac{1}{3} \int_0^1 dx_B dx_K \int_0^{1/\Lambda} b_B db_B b_K db_K \phi_B(x_B) \phi_{K^*}(x_K) \\ &\times \left\{ (1 - r^2 + x_B)(rx_K + x_B) H_8^{(a)}(Ab_K, B_8 b_B, B_8 b_K) F_8(t_{8a}) \right. \\ &\quad + [(2 - 3r)x_K - x_B + r(1 - x_K)(rx_K - 2rx_B + 3x_B)] \\ &\quad \times H_8^{(b)}(Ab_B, C_8 b_B, C_8 b_K) F_8(t_{8b}) \\ &\quad + (1 + r)(1 - r^2)[(1 + r)x_B - rx_K] \\ &\quad \times H_8^{(c)}(\sqrt{|A'^2|} b_K, D_8 b_B, D_8 b_K) F_8(t_{8c}) \\ &\quad - [(1 - r^2)((1 - r^2)(2 - x_K) + (1 + 3r)(2x_K - x_B)) \\ &\quad + 2r^2 x_K(x_K - x_B)] \\ &\quad \left. \times H_8^{(d)}(\sqrt{|A'^2|} b_B, E_8 b_B, E_8 b_K) F_8(t_{8d}) \right\} \end{aligned} \quad (25)$$

with

$$\begin{aligned} A'^2 &= (1 - r^2)(x_B - x_K) M_B^2 , \quad B_8^2 = (1 - r^2 + x_B) M_B^2 , \\ C_8^2 &= (1 - x_K) M_B^2 , \quad D_8^2 = (1 - r^2) x_B M_B^2 , \quad E_8^2 = (1 - r^2) x_K M_B^2 , \\ t_{8a} &= \max(A, B_8, 1/b_B, 1/b_K) , \\ t_{8b} &= \max(A, C_8, 1/b_B, 1/b_K) , \\ t_{8c} &= \max(\sqrt{|A'^2|}, D_8, 1/b_B, 1/b_K) , \\ t_{8d} &= \max(\sqrt{|A'^2|}, E_8, 1/b_B, 1/b_K) . \end{aligned} \quad (26)$$

The function F_8 denotes the product

$$F_8(t) = \alpha_s(t)c_8(t) \exp[-S(x_B, x_K, t, b_B, b_K)] . \quad (27)$$

The hard functions

$$\begin{aligned} H_8^{(a)}(Ab_K, B_8b_B, B_8b_K) &= H_7^{(b)}(Ab_K, B_8b_B, B_8b_K) , \\ H_8^{(b)}(Ab_B, C_8b_B, C_8b_K) &= K_0(Ab_B)h'(C_8b_B, C_8b_K) \\ H_8^{(c)}(\sqrt{|A'^2|}b_K, D_8b_B, D_8b_K) &= \\ &K_0(\sqrt{|A'^2|}b_K)h(D_8b_B, D_8b_K) , \quad A'^2 \geq 0 , \\ &i\frac{\pi}{2}H_0^{(1)}(\sqrt{|A'^2|}b_K)h(D_8b_B, D_8b_K) , \quad A'^2 < 0 , \\ H_8^{(d)}(\sqrt{|A'^2|}b_B, E_8b_B, E_8b_K) &= \\ &K_0(\sqrt{|A'^2|}b_B)h'(E_8b_B, E_8b_K) , \quad A'^2 \geq 0 , \\ &i\frac{\pi}{2}H_0^{(1)}(\sqrt{|A'^2|}b_B)h'(E_8b_B, E_8b_K) , \quad A'^2 < 0 \end{aligned} \quad (28)$$

are derived from the four diagrams.

The amplitudes with the structure $i\epsilon_{\mu\rho+-}\epsilon_\gamma^\mu\epsilon_{K^*}^{*\rho}$ are

$$\begin{aligned} M_{22} &= \Gamma^{(0)}\frac{4}{3}\int_0^1\int_0^{1-x}dy\int_0^1dx_Bdx_K\int_0^{1/\Lambda}bdb\phi_B(x_B)\phi_{K^*}(x_K) \\ &\times\alpha_s(t_2)c_2(t_2)\exp[-S(x_B, x_K, t_2, b, b)] \\ &\times\left[\left((1-r)(1-r^2)+2r^2x_K+2x_B\right)y\right. \\ &\left.-\left(r(1+r)x_K+(3-r)x_B\right)(1-x)\right] \\ &\times\frac{(1-r^2)x_Kx}{xy(1-r^2)x_KM_B^2-M_c^2}H_2(Ab, \sqrt{|B_2^2|}b) , \end{aligned} \quad (29)$$

$$M_{27} = -M_{17} , \quad (30)$$

$$\begin{aligned} M_{28} &= \Gamma^{(0)}\frac{1}{3}\int_0^1dx_Bdx_K\int_0^{1/\Lambda}b_Bdb_Bb_Kdb_K\phi_B(x_B)\phi_{K^*}(x_K) \\ &\times\left\{(1-r^2+x_B)(rx_K+x_B)H_8^{(a)}(Ab_K, B_8b_B, B_8b_K)F_8(t_{8a})\right. \\ &\left.+[(2-3r)x_K-x_B-r(1-x_K)(rx_K-2rx_B+3x_B)]\right. \\ &\left.\times H_8^{(b)}(Ab_B, C_8b_B, C_8b_K)F_8(t_{8b})\right\} \end{aligned}$$

$$\begin{aligned}
& + (1-r)(1-r^2)[(1+r)x_B - rx_K] \\
& \times H_8^{(c)}(\sqrt{|A'^2|}b_K, D_8b_B, D_8b_K)F(t_{8c}) \\
& - [(1-r^2)((1-r^2)(2+x_K) - (1-3r)x_B) \\
& - 2r^2x_K(x_K - x_B)] \\
& \times H_8^{(d)}(\sqrt{|A'^2|}b_B, E_8b_B, E_8b_K)F(t_{8d}) \Big\} . \tag{31}
\end{aligned}$$

It is obvious that the above factorization formulas bear the features of Eq. (5).

The exponentials $\exp(-S)$ appearing in M_{ij} are the complete Sudakov factor, with the exponent

$$\begin{aligned}
S = & s(x_BP_1^-, b_B) + 2 \int_{1/b_B}^t \frac{d\mu}{\mu} \gamma_q(\alpha_s(\mu)) \\
& + s(x_KP_2^+, b_K) + s((1-x_K)P_2^+, b_K) + 2 \int_{1/b_K}^t \frac{d\mu}{\mu} \gamma_q(\alpha_s(\mu)). \tag{32}
\end{aligned}$$

The wave functions ϕ_B and ϕ_{K^*} satisfy the normalization,

$$\int_0^1 \phi_i(x) dx = \frac{f_i}{2\sqrt{6}}, \tag{33}$$

with the decay constant f_i , $i = B$ and K^* . The wave function for the K^* meson with transverse polarizations has been derived using QCD sum rules [7], given by

$$\phi_{K^*} = \sqrt{6}f_{K^*}x(1-x)[0.7 - (1-2x)^2]. \tag{34}$$

As to the B meson wave functions, we investigate two models [15, 16],

$$\phi_B^{(I)}(x) = \frac{N_B x(1-x)^2}{M_B^2 + C_B(1-x)}, \tag{35}$$

$$\phi_B^{(II)}(x) = N'_B \sqrt{x(1-x)} \exp \left[-\frac{1}{2} \left(\frac{xM_B}{\omega} \right)^2 \right], \tag{36}$$

with the normalization constants N_B and N'_B , and the shape parameters C_B and ω .

4. Results

In the evaluation of the various form factors and amplitudes, we adopt $G_F = 1.16639 \times 10^{-5} \text{ GeV}^{-2}$, the decay constants $f_B = 200 \text{ MeV}$ and $f_{K^*} = 220 \text{ MeV}$, the CKM matrix elements $|V_{ts}^* V_{tb}| = 0.04$, the masses $M_c = 1.5 \text{ GeV}$, $M_B = 5.28 \text{ GeV}$ and $M_{K^*} = 0.892 \text{ GeV}$, and the $\bar{B}^0$ meson lifetime $\tau_{B^0} = 1.53 \text{ ps}$ [11]. It turns that no matter what value of the shape parameter C_B is chosen, the model $\phi_B^{(I)}$ in Eq. (35) with a flat profile leads to results much smaller than the CLEO data of the branching ratio $\mathcal{B}(B \rightarrow K^* \gamma) = (4.2 \pm 0.8 \pm 0.6) \times 10^{-5}$ [17]: the maximal prediction from $\phi_B^{(I)}$, corresponding to the shape parameter $C_B = -M_B^2$, is about 1.1×10^{-5} . The model $\phi_B^{(II)}$ in Eq. (36) with a sharp peak at small x then serves a possible candidate. It is indeed found that as $\omega = 0.425 \text{ GeV}$, a prediction 4.19×10^{-5} for the branching ratio is reached, which is equal to the central value of the experimental data. If varying the shape parameter to $\omega = 0.42 \text{ GeV}$ and to $\omega = 0.43 \text{ GeV}$, we obtain the branching ratios 4.36×10^{-5} and 4.08×10^{-5} , respectively. It indicates that the allowed range of ω is wide due to the large uncertainties of the data.

The detailed contribution from each amplitude M_{ij} is listed in the following table in the unit of 10^{-6} GeV^{-2} :

M_{12}/Γ_0	M_{17}/Γ_0	M_{18}/Γ_0
$-6.20 - 20.38i$	$11.10 - 198.08i$	$-1.94 - 0.29i$
M_{22}/Γ_0	M_{27}/Γ_0	M_{28}/Γ_0
$-4.84 - 16.17i$	$-11.10 + 198.08i$	$2.13 + 0.36i$

It is observed that the contributions from the operator O_2 are comparable with those from the penguin operator O_7 . The amplitudes associated with O_8 are less important. The other operators give contributions smaller than those from O_8 by an order of magnitude, which are thus not included here. Furthermore, the imaginary parts of $M_{1(2)7}$ from the timelike interal quarks play an essential role for the explanation of the data.

In this work we have determined the B meson wave function,

$$\phi_B(x) = 1.81359 \sqrt{x(1-x)} \exp \left[-\frac{1}{2} \left(\frac{x M_B}{0.425 \text{ GeV}} \right)^2 \right], \quad (37)$$

from the best fit to the experimental data of $\mathcal{B}(B \rightarrow K^* \gamma)$. We stress that, however, Eq. (37) is not conclusive because of the large allowed range of the shape parameter ω . A more precise B meson wave function can be obtained

by considering a global fit to the data of various decay modes, including $B \rightarrow D^{(*)}\pi(\rho)$ [18]. Once the B meson wave function is fixed, we shall employ it in the evaluation of the nonleptonic charmless decays.

This work was supported by the National Science Council of Republic of China under the Grant No. NSC-88-2112-M006-013.

References

- [1] H-n. Li and H.L. Yu, Phys. Rev. Lett. **74**, 4388 (1995); Phys. Lett. B **353**, 301 (1995); Phys. Rev. D **53**, 2480 (1996).
- [2] H-n. Li, Phys. Rev. D **52**, 3958 (1995).
- [3] H-n. Li and H.L. Yu, Phys. Rev. D **53**, 4970 (1996).
- [4] H. Georgi, Phys. Lett. B **240**, 447 (1990).
- [5] M. Bauer, B. Stech and M. Wirbel, Z. Phys. C **29**, 637 (1985); C **34**, 103 (1987).
- [6] C.H. Chang and H-n. Li, Phys. Rev. D **55**, 5577 (1997).
- [7] V.L. Chernyak and A.R. Zhitnitsky, Phys. Rep. **112**, 173 (1984).
- [8] M. Gourdin, A.N. Kamal, and X.Y. Pham, Phys. Rev. Lett. **73**, 3355 (1994).
- [9] C.E. Carlson and J. Milana, Phys. Rev. D **49**, 5908 (1994).
- [10] C.Y. Wu, T.W. Yeh, and H-n. Li, Phys. Rev. D **55**, 237 (1997).
- [11] T.W. Yeh and H-n. Li, Phys. Rev. D **56**, 1615 (1997).
- [12] N. Isgur and M.B. Wise, Phys. Lett. B **232**, 113 (1989); **237**, 527 (1990).
- [13] H-n. Li and G.L. Lin, in preparation.
- [14] A.J. Buras *et al.*, Nucl. Phys. **B424**, 374 (1994).
- [15] R. Akhoury, G. Sterman and Y.-P Yao, Phys. Rev. D **50**, 358 (1994).
- [16] M. Bauer and M. Wirbel, Z. Phys. C **42**, 671 (1989).
- [17] S. Stone, to appear in Proceedings of the Fourth Workshop on Particle Physics Phenomenology, Kaohsiung, Taiwan, 1998.
- [18] H-n. Li and B. Melic, in preparation.